

The Use of Lasers for Periodontal Debridement: Marketing Tool or Proven Therapy?

Rachel Chanthaboury, RDH, BDS; Tassos Irinakis, DDS, Dip Perio, MSc, FRCD(C)

Auteur-ressource

Dr Irinakis
Courriel : irinakis@interchange.ubc.ca



SOMMAIRE

L'emploi des lasers en dentisterie a récemment fait l'objet d'une grande attention, à la fois en pratique clinique et en recherche. Les propriétés uniques des lasers donnent des résultats cliniques désirables dans certains cas et favorisent l'acceptation des traitements par les patients. Plusieurs types de lasers utilisés en complément thérapeutique parodontal ont fait l'objet de recherches. Parmi eux, on compte les lasers au dioxyde de carbone (CO₂), les lasers à diode, les lasers au néodyme et à grenat d'yttrium-aluminium (laser Nd:YAG) et les lasers à erbium et à grenat d'yttrium-aluminium (laser Er:YAG). Toutefois, certains résultats indésirables ont été associés à chacun des types de lasers, y compris les dommages d'origine thermique aux surfaces radiculaires, l'augmentation de la température pulpaire et la production de sous-produits toxiques. Le laser Er:YAG est celui qui a donné les résultats les plus prometteurs, puisqu'il peut effectuer le débridement efficacement avec un minimum d'effets indésirables. Il faudra cependant effectuer davantage de recherches pour déterminer les méthodes et les milieux idéaux pour l'utilisation sécuritaire et efficace des lasers en pratique clinique.

Mots clés MeSH : debridement/methods; laser surgery/methods; periodontal diseases

© J Can Dent Assoc 2005; 71(9):653-8
Cet article a été révisé par des pairs.

The concept of lasers dates back to 1917 with Einstein's theory of stimulated emission, but it was not until 1960 that the first working laser was created by Theodore Maiman.¹ Lasers are currently used in a wide range of medical and cosmetic procedures including cataract surgery² and hair removal.³ However, they have only recently received attention in clinical dental settings. Lasers are being recognized for their ability to ablate hard tissues with minimal anesthesia,^{4,5} reduce bacteria counts in root canals^{6,7} and even provide hemostasis of soft tissues during their use.⁸ The aim of this article is to review various studies of lasers used in periodontal debridement.

Lasers and Clinical Dentistry

The word "laser" is an acronym for "light amplification by stimulated emission of radiation." Lasers are categorized according to the

medium used to provide atoms to the emitting system. Each type of atom can absorb photons of only certain wavelengths; therefore, each medium will produce a laser beam with a specific range of wavelengths.⁹ Light of different wavelengths will interact differently with tissue and has different adsorption qualities. Lasers used in dentistry emit wavelengths between 377 nm and 10.6 µm. The most common types are carbon dioxide (CO₂), diode, neodymium:yttrium-aluminium-garnet (Nd:YAG) and erbium:yttrium-aluminium-garnet (Er:YAG) lasers. They are currently used for cavity preparation,⁴ tooth whitening,¹⁰ gingival incisions¹¹ and other applications. In periodontics, lasers have been investigated for use in promoting periodontal attachment,¹² elimination of bacteria from periodontal pockets,⁶ debridement of root surfaces¹³ and treatment of dentinal hypersensitivity.¹⁴




Figure 1: Absorption coefficients and wavelengths of various lasers. Note the significant differences between short and long wavelength lasers. Reprinted with permission from the Institute of Laser Dentistry.

Carbon Dioxide Laser

The CO₂ laser has the longest wavelength of all dental lasers (10.6 μm). Compared with other lasers, such as the Er:YAG, its absorption coefficient in hydroxyapatite is very high (10⁴), but the absorption coefficient in water is low (10³) (Fig. 1),¹⁵ accounting for the powerful ablative properties of this laser.

Almost all studies of CO₂ lasers for periodontal debridement are in vitro; information from clinical settings is limited. All in vitro studies noted similar effects on the root surfaces of extracted teeth, i.e., charring, cratering, carbonization, surface cracking, meltdown and resolidification of minerals like calculus.¹⁶ The use of a coolant had no effect.^{15,17,18} Less aggressive results have been reported when a pulsed defocused beam was used, but this resulted in non-favourable morphologic changes in the tooth surface.¹⁹

One of the problems associated with the use of a CO₂ laser is the estimated depth of laser energy penetration into the calculus masses. In vivo, microbial plaque thickens to 100 μm within hours of colonization. Complete eradication of such a thick layer would require a penetrating wavelength with higher energy density or longer and more repeated exposures, which would lead to increased risk of damage to the adjacent tissues.¹⁶

There is also evidence that CO₂ lasers may produce toxic substances by photothermal vaporization. The high absorption of CO₂ laser energy by hydroxyapatite results in transformation of some of the radiant energy into heat, increasing the local temperature to over 700°C, which is sufficient to melt hydroxyapatite, and resulting in the emission of cyanate and cyanamide — both toxic substances.¹⁵

In vivo, CO₂ laser treatment had no significant long-term effect on levels of *Porphyromonas gingivalis*, interleukin-1 beta (IL-1β) or gingival crevicular fluid (GCF) and actually led to an increase in IL-1β.²⁰ In this study, probing

depths at 12 weeks had decreased after CO₂ laser treatment although less than that observed after Nd:YAG laser treatment and ultrasonic scaling. Improvements in clinical attachment levels were only significant in the latter 2 groups.

Diode Laser

Surgical diode lasers emit coherent, monochromatic light of short wavelength. The shorter wavelength results in a small absorption coefficient in water and hydroxyapatite; therefore, diode lasers do not ablate hard tissues very well due to their weak interaction with mineralized structures. However, this wavelength results in a high absorption coefficient in dark media such as hemoglobin, making diode lasers excellent for surgical cutting in well-vascularized soft tissues.²¹ The procedure results in coagulation and minimal bleeding when used on gingiva.

In vitro studies testing the efficiency of the diode laser in periodontal debridement have been carried out using several models: the 665-nm AlGaAs (aluminium-germanium-arsenide) laser,²² the 810-nm GaAlAs (gallium-aluminium-arsenide) laser,^{22,23} the 655-nm GaAlAs laser¹² and a 980-nm diode laser.²⁴ In all of these studies, there were minimal thermal increases over the 5°C accepted limit. The bactericidal effect of these lasers depended on species of bacterium, wavelength and dose.¹² The in vitro removal of calculus using a diode laser seemed to be consistent and comparable with manual scaling and root planing (SRP).^{12,22–24}

In in vivo studies with diode lasers, large amounts of calculus remained after treatment and there was significant structural damage to root surfaces.¹² Yilmaz and others²⁵ used a GaAlAs laser in a randomized controlled trial and found no beneficial effect over SRP alone. The differences between in vitro and in vivo studies may be attributed to the presence of blood in vivo, which may influence the amount of fluorescence radiation reaching the calculus.

Although diode lasers are currently being investigated in other aspects of dentistry, including treatment of peri-implantitis, they do not seem to provide any advantages over SRP in periodontal debridement. However, as this is a soft tissue laser, it may have potential as an adjunct to classical periodontal debridement rather than as monotherapy. Romanos and others²⁴ found the 980-nm laser to be superior to SRP in removing thin pocket epithelium. More research is needed in this area.

Neodymium:Yttrium–Aluminium–Garnet Laser

Nd:YAG lasers have a slightly longer wavelength (1,064 nm) than diode lasers. Although their absorption by healthy human enamel is very low, their much greater absorption by carious enamel makes them worthy of investigation.²⁶ The Nd:YAG laser has been tested in periodontal therapy since the early 1990s.^{27–29}

Table 1 Summary of in vitro studies of the erbium:yttrium-aluminium-garnet laser

Authors	Outcome measures	Sample size	Water coolant	Controls	Random	Pulses/s	Tip size	Energy output; mJ/pulse	Results
Aoki and others ⁴²	SEM observation of effect of laser in calculus removal SEM observation of root and thermal effects of laser with and without water coolant	53	Yes Yes	No	Yes	10 (200 µs)	600-µm diameter	10, 20, 30, 50, 80, 120	Ablated calculus and underlying cementum and dentin Maximum calculus selectivity at 30 mJ/pulse Fewer thermal effects with water irrigation Minimal rises in pulpal temperature
Gašpirc and others ³²	Morphology, chemical structure and diffusion processes of root surface	60	No	Yes	Yes	10 (250 µs)	—	60, 80, 100	Cementum and dentin removed without thermal side effects Chemical structure of root surface maintained Diffusion process increased more than with Nd:YAG laser
Frentzen and others ⁴¹	SEM and light microscopic examination of root surface characteristics and calculus removal	40	Yes	Yes	Yes	—	1.10 × 5, 1.65 × 5 mm	160	Adequate calculus removal, roughened surfaces and increased cementum removal compared with SRP Crater depths within range of those with SRP and ultrasonics
Aoki and others ⁴⁷	Histologic/SEM observation of root surface and efficiency of scaling Thermal changes	53	Yes	Yes	Yes	10 (200 µs)	600-µm diameter	40	Lower level of vibration than with ultrasonic debridement No major thermal damage Rougher root appearance than with ultrasonic debridement Settings not selective enough for calculus removal without root damage
Folwaczny and others ⁴³	SEM examination of root surface Evaluation of substance removal (calculus and root surface)	50	No	Yes	Yes	15 (250 ns)	1.65 × 0.5 mm	60, 80, 100, 150	Significantly more substance removal than controls at all energy levels No residual deposits or smear layer formation and no thermal side-effects

Nd:YAG = neodymium:yttrium-aluminium-garnet; SEM = scanning electron microscopy; SRP = scaling and root planing.

Several in vitro studies have used scanning electron microscopy to investigate the effects of Nd:YAG lasers on root surfaces.^{17,30–32} Most of these studies used a pulsed laser beam with a contact optical fibre, except for that of Wilder-Smith and colleagues,³¹ who positioned the tip of the laser 5 mm from the root surface. All 4 studies found that the Nd:YAG laser had a detrimental effect on the root surface; damage ranged from heat cracking to charring, cementum meltdown and crater formation. Although surface cooling with water, air or both decreased the damage, a 25% to 40% increase in laser energy density was needed to produce equivalent ablation effects.¹⁷

Cytokines, such as IL-1β, have been implicated in the pathogenesis of various forms of periodontal disease. Endotoxin seems to be the most potent stimulator of IL-1β production. Therefore, it seems only natural to target the removal of cementum-bound endotoxin during root planing. Liu and others³³ examined the effect of Nd:YAG lasers on endotoxin in teeth extracted due to periodontal disease. They found that the laser alone could not produce a high enough temperature to destroy cementum-bound endotoxin and still remain within safe clinical limits. SRP alone or combined with laser treatment produced similar outcomes in terms of the reduction of IL-1β for up to 3

months.³⁴ In a comparison of Nd:YAG laser monotherapy in vivo versus ultrasonic scaling, probing depth reduction and attachment levels observed at 3 months were also similar.²⁰

After observing positive results, investigators^{8,35} suggested that perhaps the Nd:YAG laser should be used as an adjunct to SRP to perform sulcular debridement rather than to remove calculus. Its role would focus on the elimination of bacterially infested pocket epithelium rather than actual calculus ablation. As pocket epithelium in a diseased environment consists of significant granulation tissue (which is a dark colour due to dilated blood vessels), the laser could stimulate evaporation of the darker tissue thus reducing periodontal pathogens and prolonging their recolonization time. The tip of the optical fibre would be brushed over the treatment area allowing light contact with the pocket epithelium for the necessary amount of time. After treating and monitoring 744 periodontal pockets (pocket depth > 4 mm), Neill and Mellonig³⁶ found that SRP alone and SRP plus Nd:YAG laser treatment produced a pocket depth reduction of 1.6 mm and 1.7 mm, respectively, which was maintained at 3 months. Clinical attachment levels and microbial counts also showed similar improvement with the 2 methods; a tendency toward better results with the combined therapy was not statistically significant.

Table 2 Summary of in vivo studies of the erbium:yttrium-aluminium-garnet laser

Authors	Outcome measures	Sample size	Water coolant	Controls	Random	Tip size mm	Energy output; mJ/pulse	Time	Conclusions
Schwarz and others ¹⁴	Clinical measurements Microbiological evaluation Follow-up at 6 months	20 patients 660 sites	Yes	Yes	Yes	1.65 × 0.5 1.10 × 0.5	160	Avg. 5 minutes/ single-rooted tooth; 10 minutes/ tooth	Significantly smaller increase in gingival recession and greater reduction in BOP and CAL gain over SRP Comparable reduction in motile rods and spirochetes to SRP
Schwarz and others ⁴⁸	Clinical measurements Microbiological evaluation Follow-up for 12 months	20 patients 600 sites	Yes	Yes	Yes	1.65 × 0.5	160	Avg. 10 minutes/ single-rooted tooth; 10 minutes/ tooth	Comparable improvements in clinical measurements and reduction in motile rods and spirochetes in both groups SRP combined with laser did not seem to improve clinical outcome over laser alone
Schwarz and others ⁴⁹	CAL gain Follow-up for 2 years	20 patients 660 sites	Yes	Yes	Yes	1.65 × 0.5 1.10 × 0.5	160	Avg. 5 minutes/ single-rooted tooth; 16 minutes/ tooth	At 1 and 2 years, laser had a significantly higher BOP reduction and CAL gain At 1 and 2 years, bacterial counts similar to baseline for both groups CAL gain in both groups maintained over 2 years

BOP = bleeding on probing; CAL = clinical attachment level; SRP = scaling and root planing.

Because clinical results are not improved by adding laser treatment to the gold standard SRP, the use of lasers as an adjunct to debridement should be questioned. Their use potentially adds to treatment time and increases the cost significantly. In a split-mouth study, Sjostrom and Friskopp⁸ investigated several parameters, including patient satisfaction and time added to treatment when an Nd:YAG laser was used as an adjunct to SRP. At 4 months, there tended to be greater improvement after laser-assisted periodontal debridement, although the difference between the groups was not significant. The authors made several important observations. They found that the use of the laser before debridement weakened the attachment of calculus to the root surface, thus facilitating its removal with hand instruments; however, the time needed for debridement increased by 15%. They also noted an analgesic effect of Nd:YAG leading to a significant decrease in requests for local anesthesia by the patients. The hemostasis effect may also prove important in patients who are on anticoagulants or have a bleeding disorder. Patients reported significantly less postoperative pain and swelling when laser treatment was used as an adjunct, although this could be due to the placebo effect that a “new” and “advanced” treatment may have on their perception.

Erbium:Yttrium–Aluminium–Garnet Laser

This is the type of laser most commonly studied for use in periodontal debridement. It has received an enormous amount of attention in dentistry since its first application in the late 1980s.^{37,38} It has a wavelength of 2,940 nm (2.94 μm), and very high absorption coefficients in water and hydroxya-

patite compared with the diode and Nd:YAG lasers. Because of its high rate of absorption in water, the Er:YAG laser ablates hard tissue through “microexplosions” rather than heating the tissue, resulting in minimal thermal side-effects.³² This desirable property of the Er:YAG laser led to its approval in 1997 by the Food and Drug Administration in the United States for use on hard tissues; it was the first dental laser to receive approval for preparation of dental cavities.³⁹ Because calculus is also a hard mineralized substance, various studies have tested the Er:YAG laser for debridement of root surfaces.

Variation in experimental design and laser settings make studies difficult to compare. Most well-designed studies in the peer-reviewed literature are in vitro. Earlier studies⁴⁰ showed detrimental outcomes of Er:YAG laser treatment on cementum surfaces of extracted teeth, including cratering and heat cracking, regardless of the presence of water coolant. Others⁴¹ noted a greater loss of cementum and residual root roughness when root planing with an Er:YAG laser compared with ultrasonics and hand instrumentation; under a scanning electron microscope, complete removal of cementum was apparent in 22.5% of laser-treated teeth compared with 12.5% of teeth treated by scaling alone. These negative outcomes were confirmed in vitro when energy outputs exceeded 50 mJ/pulse,⁴² although undesirable thermal effects were decreased in vitro when water irrigation was used.¹⁸ A water spray may decrease pulpal wall temperature by 2.2°C.³⁵ It seems that even at a relatively low radiation energy (e.g., 60 mJ), the Er:YAG laser removed considerable calculus, cementum or both, and penetration into the cementum increased significantly at 150 mJ.⁴³ Hand instrumentation

may lead to removal of up to 264.4 μm of cementum⁴⁴ or even 343.3 μm .⁴⁵ Ultrasonic instrumentation may penetrate up to 83.3 μm of cementum depending on the roughness of the diamond coating and the application force used by the investigator.⁴⁶ Up to 386.12 μm of cementum was removed from teeth with a laser at 100 mJ.⁴³ However, Aoki and others⁴⁷ disagreed with these findings and reported the maximum removal of cementum by laser of 140 μm (Table 1).

In the few in vivo studies of the Er:YAG laser (Table 2), it was used alone or in conjunction with SRP. In a comparison of laser debridement as monotherapy (160 mJ) with laser debridement as an adjunct to SRP, most clinical parameters improved similarly and remained constant over one year.⁴⁸ However, the recession readings tended to favour the monotherapy laser group, although they were not statistically significant. The authors concluded that there was no clinical benefit derived from adding SRP to laser treatment alone. The same group of investigators¹⁴ compared laser treatment with SRP as monotherapies and again obtained favourable results in the laser-treated teeth. In terms of probing depth and attachment level, both groups showed statistically significant improvement after treatment. For clinical attachment level, more improvement occurred in the laser-treated group than the SRP group, but the difference was not statistically significant. A 2-year prospective study⁴⁹ by the same investigator group confirmed the favourable results with lasers. Sculean and others⁵⁰ compared the use of a laser alone with ultrasonic debridement and found the 2 treatments comparable in effectiveness of calculus removal, although the laser was less efficient than ultrasonic debridement. No differences in clinical improvement were noted.

More prospective studies are necessary to confirm the optimistic results offered by these studies. The Er:YAG laser seems to have a place in periodontal debridement, but it remains to be seen whether predictably positive outcomes can be achieved using it as an adjunct or as a monotherapy.

Conclusions

Bacterial invasion cannot be eradicated by mechanical debridement alone. In some cases, combining mechanical therapy with laser treatment appears beneficial. Some recent controlled studies favour laser-assisted periodontal debridement with a soft-tissue laser or use of a hard-tissue laser as a monotherapy. More in vivo studies are needed to establish the safety and benefit of dental lasers, either as an adjunct or as an alternative to the traditional methods of scaling and root planing. ➔

THE AUTHORS



Ms. Chanthaboury is a full-time registered dental hygienist in private practice in Vancouver, B.C.



Dr. Irinakis is a full-time clinical associate professor and director of graduate periodontics and implant surgery at the University of British Columbia, Vancouver, B.C. He is also in part-time private practice in periodontics in Vancouver.

Correspondence to: Dr. Tassos Irinakis, Faculty of Dentistry, University of British Columbia, 2199 Wesbrook Mall, Vancouver, BC V6T 1Z3. E-mail: irinakis@interchange.ubc.ca.

The authors have no declared financial interests in any company manufacturing the types of products mentioned in this article.

References

- Maiman TH. Stimulated optical radiation in ruby. *Nature* 1960; 187:493–4.
- Verges C, Llevat E. Laser cataract surgery: technique and clinical results. *J Cataract Refract Surg* 2003; 29(7):1339–45.
- Bouzari N, Tabatabai H, Abbasi Z, Firooz A, Dowlati Y. Laser hair removal: comparison of long-pulsed Nd:YAG, long-pulsed alexandrite, and long-pulsed diode lasers. *Dermatol Surg* 2004; 30(4 Pt 1):498–502.
- Oelgiesser D, Blasbalg J, Ben-Amar A. Cavity preparation by Er-YAG laser on pulpal temperature rise. *Am J Dent* 2003; 16(2):96–8.
- Jayawardena JA, Kato J, Moriya K, Takagi Y. Pulpal response to exposure with Er:YAG laser. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2001; 91(2):222–9.
- Ando Y, Aoki A, Watanabe H, Ishikawa I. Bactericidal effect of erbium YAG laser on periodontopathic bacteria. *Lasers Surg Med* 1996; 19(2):190–200.
- Meral G, Tasar F, Kocagoz S, Sener C. Factors affecting the antibacterial effects of Nd:YAG laser in vivo. *Lasers Surg Med* 2003; 32(3):197–202.
- Sjostrom L, Friskopp J. Laser treatment as an adjunct to debridement of periodontal pockets. *Swed Dent J* 2002; 26(2):51–7.
- Miserendino LJ, Neiburger EJ, Pick RM. Current status of lasers in dentistry. *Ill Dent J* 1987; 56(4):254–7.
- Luk K, Tam L, Hubert M. Effect of light energy on peroxide tooth bleaching. *J Am Dent Assoc* 2004; 135(2):194–201.
- Neiburger EJ. Rapid healing of gingival incisions by the helium-neon diode laser. *J Mass Dent Soc* 1999; 48(1):8–13, 40.
- Kreisler M, Meyer C, Stender E, Daublander M, Willershausen-Zonnchen B, d'Hoedt B. Effect of diode laser irradiation on the attachment rate of periodontal ligament cells: an in vitro study. *J Periodontol* 2001; 72(10):1312–7.
- Schwarz F, Sculean A, Berakdar M, Szathmari L, Georg T, Becker J. In vivo and in vitro effects of an Er:YAG laser, a GaAlAs diode laser, and scaling and root planing on periodontally diseased root surfaces: a comparative histologic study. *Lasers Surg Med* 2003; 32(5):359–66.
- Schwarz F, Sculean A, Georg T, Reich E. Periodontal treatment with an Er:YAG laser compared to scaling and root planing. A controlled clinical study. *J Periodontol* 2001; 72(3):361–7.
- Sasaki KM, Aoki A, Masuno H, Ichinose S, Yamada S, Ishikawa I. Compositional analysis of root cementum and dentin after Er:YAG laser irradiation compared with CO₂ laser and intact roots using Fourier transformed infrared spectroscopy. *J Periodontol Res* 2002; 37(1):50–9.
- Tucker D, Cobb CM, Rapley JW, Killoy WJ. Morphologic changes following in vitro CO₂ laser treatment of calculus-laden root surfaces. *Lasers Surg Med* 1996; 18(2):150–6.
- Israel M, Cobb CM, Rossmann JA, Spencer P. The effects of CO₂, Nd:YAG and Er:YAG lasers with and without surface coolant on tooth root surfaces. An in vitro study. *J Clin Periodontol* 1997; 24(9 Pt 1):595–602.
- Sasaki KM, Aoki A, Ichinose S, Ishikawa I. Morphological analysis of cementum and root dentin after Er:YAG laser irradiation. *Lasers Surg Med* 2002; 31(2):79–85.
- Barone A, Covani U, Crespi R, Romanos GE. Root surface morphological changes after focused versus defocused CO₂ laser irradiation: a scanning electron microscopy analysis. *J Periodontol* 2002; 73(4):370–3.
- Miyazaki A, Yamaguchi T, Nishikata J, Okuda K, Suda S, Orima K, and others. Effects of Nd:YAG and CO₂ laser treatment and ultrasonic scaling on periodontal pockets of chronic periodontitis patients. *J Periodontol* 2003; 74(2):175–80.
- Bach G, Neckel C, Mall C, Krekeler G. Conventional versus laser-assisted therapy of periimplantitis: a five-year comparative study. *Implant Dent* 2000; 9(3):247–51.

22. Chan Y, Lai CH. Bactericidal effects of different laser wavelengths on periodontopathic germs in photodynamic therapy. *Lasers Med Sci* 2003; 18(1):51–5.
23. Theodoro LH, Haypek P, Bachmann L, Garcia VG, Sampaio JEC, Zezell DM, and other. Effect of Er:YAG and diode laser irradiation on the root surface: morphological and thermal analysis. *J Periodontol* 2003; 74(6):838–43.
24. Romanos GE, Henze M, Banihashemi S, Parsanejad HR, Winckler J, Nentwig GH. Removal of epithelium in periodontal pockets following diode (980 nm) laser application in the animal model: an in vitro study. *Photomed Laser Surg* 2004; 22(3):177–83.
25. Yilmaz S, Kuru B, Kuru L, Noyan U, Argun D, Kadir T. Effect of gallium arsenide diode laser on human periodontal disease: a microbiological and clinical study. *Lasers Surg Med* 2002; 30(1):60–6.
26. Harris J, Walmsley D. Technology transfer in dentistry: a network resource for clinicians, scientists, manufacturers and patients. *Dent Update* 2002; 29(7):338–9.
27. Morlock BJ, Pippin DJ, Cobb CM, Killoy WJ, Rapley JW. The effect of Nd:YAG laser exposure on root surfaces when used as an adjunct to root planing: an in vitro study. *J Periodontol* 1992; 63(7):637–41.
28. Cobb CM, McCawley TK, Killoy WJ. A preliminary study on the effects of the Nd:YAG laser on root surfaces and subgingival microflora in vivo. *J Periodontol* 1992; 63(8):701–7.
29. Ito K, Nishikata J, Murai S. Effects of Nd:YAG laser radiation on removal of a root surface smear layer after root planing: a scanning electron microscopic study. *J Periodontol* 1993; 64(6):547–52.
30. Thomas D, Rapley J, Cobb C, Spencer P, Killoy W. Effects of the Nd:YAG laser and combined treatments on in vitro fibroblast attachment to root surfaces. *J Clin Periodontol* 1994; 21(1):38–44.
31. Wilder-Smith P, Arrastia AM, Schell MJ, Liaw LH, Grill G, Berns MW. Effect of Nd:YAG laser irradiation and root planing on the root surface: structural and thermal effects. *J Periodontol* 1995; 66(12):1032–9.
32. Gaspirc B, Skaleric U. Morphology, chemical structure and diffusion processes of root surface after Er:YAG and Nd:YAG laser irradiation. *J Clin Periodontol* 2001; 28(6):508–16.
33. Liu CM, Shyu YC, Pei SC, Lan WH, Hou LT. In vitro effect of laser irradiation on cementum-bound endotoxin isolated from periodontally diseased roots. *J Periodontol* 2002; 73(11):1260–6.
34. Liu CM, Hou LT, Wong MY, Lan WH. Comparison of Nd:YAG laser versus scaling and root planing in periodontal therapy. *J Periodontol* 1999; 70(11):1276–82.
35. Radvar M, MacFarlane TW, MacKenzie D, Whitters CJ, Payne AP, Kinane DF. An evaluation of the Nd:YAG laser in periodontal pocket therapy. *Br Dent J* 1996; 180(2):57–62.
36. Neill ME, Mellonig JT. Clinical efficacy of the Nd:YAG laser for combination periodontitis therapy. *Pract Periodontics Aesthet Dent* 1997; 9(6 Suppl):1–5.
37. Hibst R, Keller U. Experimental studies of the application of the Er:YAG laser on dental hard substances: I. Measurement of the ablation rate. *Lasers Surg Med* 1989; 9(4):338–44.
38. Keller U, Hibst R. Experimental studies of the application of the Er:YAG laser on dental hard substances: II. Light microscopic and SEM investigations. *Lasers Surg Med* 1989; 9(4):345–51.
39. Cozean C, Arcoria CJ, Pelagalli J, Powell GL. Dentistry for the 21st century? Erbium:YAG laser for teeth. *J Am Dent Assoc* 1997; 128(8):1080–87.
40. Dostalova T, Jelinkova H, Krejsa O, Hamal H. Evaluation of the surface changes in enamel and dentin due to possibility of thermal overheating induced by Erbium:YAG laser radiation. *Scanning Microsc* 1996; 10(1):285–90.
41. Frentzen M, Braun A, Aniol D. Er:YAG laser scaling of diseased root surfaces. *J Periodontol* 2002; 73(5):524–30.
42. Aoki A, Ando Y, Watanabe H, Ishikawa I. In vitro studies on laser scaling of subgingival calculus with an Erbium:YAG laser. *J Periodontol* 1994; 65(12):1097–106.
43. Folwaczny M, Mehl A, Haffner C, Benz C, Hickel R. Root substance removal with Er:YAG laser radiation at different parameters using a new delivery system. *J Periodontol* 2000; 71(2):147–55.
44. Ritz L, Hefti AF, Rateitschak KH. An in vitro investigation on the loss of root substance in scaling with various instruments. *J Clin Periodontol* 1991; 18(9):643–7.
45. Zappa U, Smith B, Simiona C, Graft H, Case D, Kim W. Root surface removal by scaling and root planing. *J Periodontol* 1991; 62(12):750–4.
46. Lavespere JE, Yukna RA, Rice DA, LeBlanc DM. Root surface removal with diamond coated ultrasonic instruments: an in vitro and SEM study. *J Periodontol* 1996; 67(12):1281–7.
47. Aoki A, Miura M, Akiyama F, Nakagawa N, Tanaka J, Oda S, and others. In vitro evaluation of Er:YAG laser scaling of subgingival calculus in comparison with ultrasonic scaling. *J Periodont Res* 2000; 35:266–77.
48. Schwarz F, Sculean A, Berakdar M, Georg T, Reich E, Becker J. Clinical evaluation of an Er:YAG laser combined with scaling and root planing for non-surgical periodontal treatment. A controlled, prospective clinical study. *J Clin Periodontol* 2003; 30(1):26–34.
49. Schwarz F, Sculean A, Berakdar M, Georg T, Reich E, Becker J. Periodontal treatment with an Er:YAG laser or scaling and root planing. A 2-year follow-up split-mouth study. *J Periodontol* 2003; 74(5):590–6.
50. Sculean A, Schwarz F, Berakdar M, Romanos GE, Arweiler NB, Becker J. Periodontal treatment with an Er:YAG laser compared to ultrasonic instrumentation: a pilot study. *J Periodontol* 2004; 75(7):966–73.