

# Upper Body Musculoskeletal Symptoms in Sardinian Dental Students

(Symptômes musculosquelettiques du haut du corps chez des étudiants  
en médecine dentaire sardes)

- Marcello Melis, DMD, RPharm •
- Youssef S. Abou-Atme, DDS, MS •
- Luca Cottogno •
- Roberta Pittau, DMD •

## S o m m a i r e

**Objet :** Le but de cette étude était de déterminer la précocité d'apparition des problèmes musculosquelettiques au cours de la profession dentaire, en comparant la prévalence des symptômes en question dans une population d'étudiants en médecine dentaire de Sardaigne (Italie), dans un échantillon témoin d'étudiants en psychologie de la même université et dans un échantillon d'étudiants en médecine dentaire du Liban.

**Méthodologie :** Cent quatorze étudiants en médecine dentaire provenant de l'Université de Cagliari (Cagliari, Sardaigne, Italie) ont été interrogés par questionnaire au sujet de la présence de plusieurs symptômes du haut du corps. Les réponses qu'ils ont données ont été comparées à celles de 2 populations différentes, soit 114 étudiants en psychologie de la même université, assortis pour l'âge et le sexe, et 178 étudiants en médecine dentaire de l'Université de Saint Joseph (Beyrouth, Liban). Les étudiants en psychologie ont été choisis pour la comparaison en raison de la dissimilitude générale entre leur activité quotidienne et celle des dentistes.

**Résultats :** Les étudiants en médecine dentaire de l'Italie et du Liban n'ont pas été différents quant à la présence des symptômes. Cependant, les étudiants en médecine dentaire de l'Italie différaient des étudiants en psychologie à un seul égard, les premiers ayant signalé davantage de douleurs au bas du dos. Le mal de tête était le symptôme le plus prévalent dans les 3 groupes et il était plus prévalent chez les femmes. Toutefois, cette différence entre les sexes n'a pas atteint la signification statistique pour les étudiants en psychologie.

**Conclusions :** L'apparition des symptômes musculosquelettiques chez les étudiants en médecine dentaire, même après une période relativement courte de formation clinique, donne à penser qu'on devrait inclure l'ergonomie dans le système d'éducation afin de réduire les risques pour les praticiens.

**Mots clés MeSH :** back pain/epidemiology; headache; muscular diseases; occupational diseases/epidemiology

© J Can Dent Assoc 2004; 70(5):306-10  
Cet article a été révisé par des pairs.

The working environment has a major role in the development of many musculoskeletal problems, although most of these disorders can be avoided or at least reduced with more attention to ergonomics.<sup>1-6</sup> In particular, dental professionals often assume positions that are uncomfortable and asymmetric, keeping the head forward and rotated to the side with the arms held out from the body. This attitude, if held for prolonged periods each day, tends to overstress muscles and joints, especially those of the neck, back

and shoulder, causing symptoms such as headache, neck and shoulder pain, and backache.<sup>1-12</sup>

Seventy-two percent of the dentists examined by Rundcrantz and others<sup>1</sup> reported some pain and discomfort of the neck, shoulders or head. In support of an occupational cause of the symptoms, pain and discomfort were less in subjects who had adopted ergonomic techniques, such as working in a standing position; altering the working position in relation to the patient, according to the specific area of the

**Table 1** Prevalence of symptoms and  $\chi^2$  analysis for comparison of dental students and psychology students

Symptom	No. (and %) of students with symptoms		$\chi^2$	<i>p</i> (df = 1)
	USC ( <i>n</i> = 114)	PSY ( <i>n</i> = 114)		
Headache	58 (50.9)	65 (57.0)	0.86	> 0.05
Neck pain	46 (40.4)	34 (29.8)	2.77	> 0.05
Lower back pain	37 (32.5)	21 (18.4)	5.92	< 0.05
Upper back pain	24 (21.1)	30 (26.3)	0.87	> 0.05
Arm symptoms	22 (19.3)	12 (10.5)	3.46	> 0.05

USC = dental students from University of Cagliari (Cagliari, Italy)

PSY = psychology students from University of Cagliari

df = degrees of freedom

mouth to be worked on; using a mirror for a better view of tooth surfaces difficult to access directly; using a wedge cushion to better position the head of the patient; and interrupting work with short pauses.<sup>1,2</sup> Similar results were reported by Kerosuo and others,<sup>12</sup> who discovered that 70% of general dental practitioners and 72% of orthodontists surveyed suffered from musculoskeletal symptoms. The sites most frequently involved were the neck (53% of those with pain) and shoulders (56% of those with pain). Chang<sup>5</sup> suggested proper selection of surgical telescopes and use of coaxial illumination headlights to prevent or eliminate chronic neck and back pain. In contrast, Marshall and others<sup>7</sup> questioned whether modifying work position or taking short pauses throughout the day would decrease symptoms.

In comparisons of dentists and subjects working in a different environment, such as farmers and pharmacists, the radiological signs and symptoms of upper body pathologic conditions were more prevalent among dentists.<sup>2,13,14</sup> Compared with office employees, dentists showed more severe symptoms; these differences were especially evident in female subjects.<sup>12</sup>

To determine how early in a dental career these musculoskeletal problems develop, the prevalence of symptoms in a dental student population in Sardinia (Italy), a control sample of psychology students from the same university (matched for age and sex) and a sample of dental students in Lebanon was determined and compared.

## Materials and Methods

Dental students at the University of Cagliari in Cagliari, Sardinia, Italy, psychology students from the same university and dental students at the University of Saint Joseph in Beirut, Lebanon, were asked to respond to a questionnaire (see Appendix I, Student survey on musculoskeletal pain, at <http://www.cda-adc.ca/jcda/vol-70/issue-5/306.html>) about the recent presence of the following body symptoms: headache, neck pain, upper back pain or stiffness, lower back pain or stiffness, and arm symptoms (pain, tingling or numbness). These symptoms were chosen because the neck and back are the body sites reportedly most affected by altered posture during dental work, and problems at those locations

**Table 2** Prevalence of symptoms and  $\chi^2$  analysis for comparison of Italian and Lebanese dental students

Symptom	No. (and %) of students with symptoms		$\chi^2$	<i>p</i> (df = 1)
	USC ( <i>n</i> = 114)	USJ ( <i>n</i> = 178)		
Headache	58 (50.9)	101 (56.7)	0.92	> 0.05
Neck pain	46 (40.4)	80 (44.9)	1.39	> 0.05
Lower back pain	37 (32.5)	66 (37.1)	1.28	> 0.05
Upper back pain	24 (21.1)	52 (29.2)	2.68	> 0.05
Arm symptoms	22 (19.3)	45 (25.3)	1.32	> 0.05

USC = dental students from University of Cagliari (Cagliari, Italy)

USJ = dental students from University of Saint Joseph (Beirut, Lebanon)

df = degrees of freedom

may lead to referred symptoms such as headache and arm symptoms.<sup>1-12</sup>

The reported symptoms were compared by means of  $\chi^2$  tests applied to  $2 \times 2$  tables. In the first analysis, the Italian dental students were compared with the psychology students (matched for age and sex); in the second analysis the Italian dental students were compared with the Lebanese dental students.

## Results

All but one of the Italian dental students returned a completed questionnaire (*n* = 114; 60 men and 54 women). The students ranged in age from 18 to 51 years (mean 22.7 years), but only one student was older than 34. Responses were obtained from 114 psychology students matched for age and sex. All of the Lebanese dental students returned the questionnaire (*n* = 178; 117 men and 61 women, mean age 20.7 years, range 17-27 years).

In the comparison of Italian dental students and psychology students (Table 1), the only difference was more lower back pain among dental students (*p* < 0.05). There were no differences between the 2 groups of dental students (*p* > 0.05; Table 2).

Headache was the most prevalent symptom in all 3 groups (Fig. 1). In a separate analysis, the prevalence of headache was compared between men and women in each student group. In all groups, a greater proportion of women than men reported this symptom (Table 3, Fig. 2), but the difference was significant only for the 2 groups of dental students (University of Cagliari:  $\chi^2 = 12.77$ , *p* < 0.01; University of St. Joseph:  $\chi^2 = 6.47$ , *p* < 0.05). In addition, the prevalence of lower back pain was higher among female psychology students than among male psychology students (*p* < 0.01; data not shown).


## Discussion

Headache was the most frequent symptom in all 3 populations examined. The percentage of subjects reporting headache (50.9% to 57.0% in the 3 groups) was higher than that reported in most other studies. Rundcrantz and others found that 38% of Swedish dentists in one study<sup>1</sup> and 52 (36.4%) of


**Table 3** Prevalence of headache in women and men in the 3 student groups

Symptom	USC (n = 114)		PSY (n = 114)		USJ (n = 178)	
	Women	Men	Women	Men	Women	Men
Headache (no. and %)	37 (68.5)	21 (35.0)	34 (63.0)	31 (51.7)	42 (68.9)	58 (49.6)
No headache (no. and %)	17 (31.5)	39 (65.0)	20 (37.0)	29 (48.3)	19 (31.1)	59 (50.4)
Total	54	60	54	60	61	117
$\chi^2$	12.77		1.15		6.47	
P	< 0.01		> 0.05		< 0.05	

USC = dental students from University of Cagliari (Cagliari, Italy), PSY = psychology students from University of Cagliari, USJ = dental students from University of Saint Joseph (Beirut, Lebanon)



**Figure 1:** Prevalence of upper body symptoms. USC = dental students from University of Cagliari (Cagliari, Italy), USJ = dental students from University of Saint Joseph (Beirut, Lebanon), PSY = psychology students from University of Cagliari. Lower back pain was significantly more prevalent within the USC group than within the PSY group ( $p < 0.05$ ).



**Figure 2:** Prevalence of headache. USC = dental students from University of Cagliari (Cagliari, Italy), USJ = dental students from University of Saint Joseph (Beirut, Lebanon), PSY = psychology students from University of Cagliari. For both groups of dental students, the difference between the sexes was statistically significant (for USC,  $\chi^2 = 12.78$ ,  $p < 0.01$ ; for USJ,  $\chi^2 = 6.47$ ,  $p < 0.05$ ).

143 dentists in another study<sup>2</sup> suffered from headache. Kerosuo and others<sup>12</sup> reported a much lower prevalence of headache (18%) among Finnish dentists and orthodontists. The mean age of subjects examined by Rundcrantz and others<sup>1,2</sup> (in their 40s) was much greater than the mean age of the students in this study (early 20s). However, Rundcrantz and others<sup>1</sup> noted that younger dentists had more headaches than older dentists. The findings of Marshall and others<sup>7</sup> are closer to the results reported here: they found a 58% prevalence of headache among Australian dentists.

Differences between men and women in terms of headache prevalence have been described,<sup>15,16</sup> although there may be variations with different types of headache.<sup>16</sup> In a questionnaire-based study conducted in Hong Kong<sup>15</sup> there was a greater preponderance of all types of headache among female subjects, regardless of age. The same study noted that clerical or service work, housekeeping, and professional or administrative jobs were the most common occupations among respondents with recurrent headaches. Studies on dental students and dental professionals,<sup>1,2,7</sup> including the one reported here, seem to agree that headaches are more common among women. However, this distinction was not evident among psychology students.

Previous studies have suggested a higher prevalence of neck pain among dentists, probably because they assume positions that don't conform to usual body positioning during work,<sup>1,2,12,17</sup> especially when compared with subjects in other occupations, where body positioning is more natural.<sup>18</sup> Although there seems to be a correlation between neck pain and occupational stance in general,<sup>19-21</sup> other studies and the results reported here do not support such a correlation for dental practitioners: the prevalence of neck pain did not differ significantly between the 2 dental student groups or between the dental students and the psychology students at Cagliari University, although there was a trend for more dental students than psychology students to report neck pain.

Rundcrantz and others<sup>17</sup> reported that 44% to 48% of men and 61% to 62% of women experienced neck pain; Niemi and others<sup>18</sup> found that only 10% of young men and 21% of young women had neck pain, but these researchers were examining different populations (dentists were surveyed in the first study, high school students were surveyed in the second study). In another study, the latter authors found that young women suffered more and sought more help with their neck and shoulder pain.<sup>22</sup> In that study, young women who participated in dynamic sports involving the upper extremities had fewer symptoms than those who participated in more static

activities. In the study reported here, there was no difference between men and women in the prevalence of neck pain.

The prevalence of upper back and shoulder pain has varied from 10% to 73% in previous studies.<sup>1,2,10,15</sup> A greater prevalence of back pain among women has been reported by some authors.<sup>12,17</sup> Lower back pain was cited as the most prevalent symptom among Australian dentists (experienced by 64% of those who suffered pain).<sup>7</sup> Given the wide range in previous results, comparison with the current findings is difficult. In a study of Swedish dentists, neither headache nor lower back pain increased over a 3-year period, and general musculoskeletal symptoms were significantly less prevalent among office workers used as controls.<sup>17</sup> The latter results are in accordance with the findings reported here; however, because we did not evaluate the symptoms longitudinally, we are unable to confirm their stability over time.

Feldman and others<sup>23</sup> and Chang<sup>5</sup> suggested that modifying risk factors for lower back pain might prevent the development of lower back pain in the future. Mendez and Gomez-Conesa<sup>24</sup> showed that programs involving practice of manoeuvres and motivating strategies impart health knowledge and habits more efficiently than those restricted to the mere transmission of information. In the authors' opinion, suitable information about appropriate arrangement of equipment and proper working postures, as well as practice in using those postures, would reduce the risk of developing musculoskeletal symptoms.

There were no significant differences in the prevalence of arm symptoms (pain, tingling, numbness) between the 2 dental student groups (19.3% among the Italian students and 25.3% among the Lebanese students) or between the Italian dental students and the psychology students (19.3% and 10.5% respectively). Hand symptoms occurred in a similar proportion of Finnish dentists (17%),<sup>12</sup> but no data were provided to allow comparison of dentists with office employees, except in terms of all locomotor symptoms combined.<sup>12</sup> In another study, elbow symptoms occurred in 13% of subjects and wrist and hand symptoms in 24% of subjects.<sup>1</sup> At 3-year follow-up of the same population,<sup>17</sup> elbow symptoms declined from 11% to 9% among male subjects and rose from 15% to 19% among female subjects; symptoms of the wrist and hand rose from 14% to 19% among male subjects and from 32% to 35% among female subjects. In the latter study<sup>17</sup> the female dentists had significantly greater symptoms than the male dentists. The questionnaire used in the current study asked about arm pain, tingling and numbness without any specific reference to hand or wrist symptoms; therefore, the results cannot be compared.

Marshall and others<sup>7</sup> reported that neurological symptoms were more common in the dominant arm of dentists than the nondominant arm; however, information of this nature was not available for the student groups assessed in this study, because no examination was performed to establish arm dominance of the subjects.

## Conclusions

In the study reported here, low back pain was the only type of upper body pain occurring significantly more frequently among dental students than among a control group of psychology students. Dental studies and dental work often involve time spent in static, uncomfortable positions, which can lead to musculoskeletal symptoms even over the relatively short clinical training period. There is a critical need to address ergonomic issues in the educational system and to change the way dentistry is practised to lower the risks to dental practitioners. ♦



*Le Dr Melis exerce dans un cabinet privé à Cagliari, en Italie.*



*Le Dr Abou-Atme exerce dans un cabinet privé à Beyrouth, au Liban.*



*M. Cottogno est étudiant en psychologie à Cagliari, en Italie.*



*La Dre Pittau exerce dans un cabinet privé à Cagliari, en Italie.*

*Écrire au : Dr Marcello Melis, Via Grosseto, 1, 09125 – Cagliari, Italie. Courriel : [marcellomelis01@tin.it](mailto:marcellomelis01@tin.it).*

*Les auteurs n'ont aucun intérêt financier déclaré.*

## Références

1. Rundcrantz BL, Johnsson B, Moritz U. Cervical pain and discomfort among dentists. Epidemiological, clinical and therapeutic aspects. Part 1. A survey of pain and discomfort. *Swed Dent J* 1990; 14(2):71–80.
2. Rundcrantz BL, Johnsson B, Moritz U. Occupational cervico-brachial disorders among dentists. Analysis of ergonomics and locomotor functions. *Swed Dent J* 1991; 15(5):105–15.
3. Laderas S, Felsenfeld AL. Ergonomics and the dental office: an overview and consideration of regulatory influences. *J Calif Dent Assoc* 2002; 30(2):135,137–8.
4. Rucker LM, Sunell S. Ergonomic risk factors associated with clinical dentistry. *J Calif Dent Assoc* 2002; 30(2):139–48.
5. Chang BJ. Ergonomic benefits of surgical telescope systems: selection guidelines. *J Calif Dent Assoc* 2002; 30(2):161–9.
6. Yoser AJ, Mito RS. Injury prevention for the practice of dentistry. *J Calif Dent Assoc* 2002; 30(2):170–6.
7. Marshall ED, Duncombe LM, Robinson RQ, Kilbreath SL. Musculoskeletal symptoms in New South Wales dentists. *Austr Dent J* 1997; 42:240–6.
8. Milerad E, Ekenvall L. Symptoms of the neck and upper extremities in dentists. *Scand J Work Environ Health* 1990; 16:129–34.
9. Shugars D, Miller D, Williams D, Fishburne C, Strickland D. Musculoskeletal pain among general dentist. *Gen Dent* 1987; 35(4):272–6.
10. Fox JG, Jones JM. Occupational stress in dental practice. *Br Dent J* 1967; 123(10):465–73.

11. Biller FE. Occupational hazards in dental practice. *Oral Hygiene* 1946; 36:1194.
12. Kerosuo E, Kerosuo H, Kanerva L. Self-reported health complaints among general dental practitioners, orthodontists, and office employees. *Acta Odontol Scand* 2000; 58(5):207-12.
13. Katevuo K, Aitasalo K. Skeletal changes in dentists and farmers in Finland. *Community Dent Oral Epidemiol* 1985; 13(1):23-5.
14. Powell M, Eccles JD. The health and work of two professional groups: dentists and pharmacists. *Dent Pract Dent Rec* 1970; 20(11):373-8.
15. Cheung RT. Prevalence of migraine, tension-type headache, and other headaches in Hong Kong. *Headache* 2000; 40(6):473-9.
16. Saper JR, Silberstein SD, Gordon CD, Hamel RL, Swidan S. Handbook of headache management. 2nd ed. Baltimore (MD): Lippincott Williams & Wilkins; 1999. p. 1-4.
17. Rundcrantz BL, Johnsson B, Moritz U. Pain and discomfort in the musculoskeletal system among dentists. A prospective study. *Swed Dent J* 1991; 15(5):219-28.
18. Niemi SM, Levoska S, Rekola KE, Keinanen-Kiukaanniemi SM. Neck and shoulder symptoms of high school students and associated psychosocial factors. *J Adolesc Health* 1997; 20(3):238-42.
19. Ehrmann-Feldman D, Shrier I, Rossignol M, Abenhaim L. Risk factors for the development of neck and upper limb pain in adolescents. *Spine* 2002; 27(5):523-8.
20. Feldman DE, Shrier I, Rossignol M, Abenhaim L. Work is a risk factor for adolescent musculoskeletal pain. *J Occup Environ Med* 2002; 44(10):956-61.
21. Szeto GP, Lee R. An ergonomic evaluation comparing desktop, notebook, and subnotebook computers. *Arch Phys Med Rehabil* 2002; 83(4):527-32.
22. Niemi SM, Levoska S, Rekola KE, Keinanen-Kiukaanniemi SM. Neck and shoulder symptoms and leisure time activities in high school students. *J Orthop Sports Phys Ther* 1996; 24(1):25-9.
23. Feldman DE, Shrier I, Rossignol M, Abenhaim L. Risk factors for the development of low back pain in adolescence. *Am J Epidemiol* 2001; 154(1):30-6.
24. Mendez FJ, Gomez-Conesa A. Postural hygiene program to prevent low back pain. *Spine* 2001; 26(11):1280-6.

**Appendix I Student survey of musculoskeletal pain**Age: \_\_\_\_\_ Gender: M  F 

Do you suffer, or have you recently suffered, from one of the following symptoms?

Put an X in the corresponding box.

		No	Mild	Moderate	Severe
Headache	Left	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
	Right	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Arm pain/tingling/numbness	Left	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
	Right	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Neck pain/stiffness	Left	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
	Right	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Upper back pain/stiffness	Left	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
	Right	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Lower back pain/stiffness	Left	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
	Right	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

Be sure you have completed every row. If you do not have the symptom described, put an X in the box corresponding with "No."

*Thank you for your time!*