

# Clinical Management of 3 Mandibular Incisors with 2 Separate Canals and Foramina : Case Report

(Prise en charge clinique de 3 incisives inférieures avec 2 canaux et un foramen distincts : rapport de cas)

- Young hee Hwang, DDS •
- Kyung san Min, DDS, MS •

## S o m m a i r e

*Le présent rapport de cas décrit le traitement radiculaire de 3 incisives inférieures avec 2 canaux et 1 foramen distinct chez un patient. Le cas montre une configuration morphologique inhabituelle du système canalaire et complète les rapports précédents sur l'existence de telles configurations canalaires dans les incisives inférieures.*

**Mots clés MeSH :** dental pulp cavity/anatomy & histology; incisor/anatomy & histology; root canal therapy

© J Can Dent Assoc 2005; 71(6):388-9  
Cet article a été révisé par des pairs.

It is generally accepted that many mandibular incisors have 2 canals, which may merge into 1 canal before reaching the apex.<sup>1-4</sup> In rare cases, separate foramina may form. In a radiographic study of 364 specimens, Benjamin and Dowson<sup>4</sup> reported that 41.4% of the mandibular incisors they studied had 2 separate canals; of these, only 1.3% had 2 separate foramina. In a study of 1,085 specimens, Miyashita and others<sup>5</sup> reported that only 3.1% of the samples had separate canals and foramina. In a study of mandibular incisors in which a surgical resection method was used to study canal anatomy, Mauger and others<sup>6</sup> demonstrated that, at the apical 1, 2 and 3 mm levels in the mandibular incisor, the canal is only rarely separated by hard tooth structure and that only 2% of the teeth they studied had 2 canals at the 1 mm resection level.

This case report describes the occurrence of 3 mandibular incisors, each with 2 separate canals and foramina, in 1 patient.

### Case Report

A 41-year-old man presented with a fractured left mandibular lateral incisor. The medical history was noncontributory, but the patient had experienced traumatic injury to the anterior mandibular region about 15 years previously, and there was a history of periodic tooth discomfort on biting. The clinical examination revealed pain on palpation and no response to thermal sensitivity tests. The mandibular central

incisors and left lateral incisor felt unusual on percussion, and the right central incisor showed severe discoloration. Preoperative radiographic examination revealed periapical radiolucent areas (Fig. 1). The right central incisor showed a radiolucent canal that stopped abruptly in the middle third of the root, indicating the presence of 2 canals (Fig. 1). The root canal, which had been partially treated, displayed a multicanal morphology (Fig. 1).

The diagnosis was pulpal necrosis with chronic apical periodontitis of the mandibular central incisors and the left lateral incisor. Root canal treatment was indicated.

After placement of a rubber dam, the deficient root canal filling material in the left lateral incisor was removed by means of Hedström files (Dentsply Maillefer, Ballaigues, Switzerland). The working lengths were estimated using an apex locator (Root ZX, Morita, Tokyo, Japan) and then confirmed by radiography (Fig. 2).

The root canals were prepared with Protaper rotary instruments (Dentsply Maillefer) and were copiously irrigated with 5.25% sodium hypochlorite. When the canal preparation was almost complete, the opening of a second root canal was noted in all 3 teeth. The working lengths were re-estimated with the apex locator and then confirmed by radiography to determine the anatomic relationship between the root canals of each tooth. The radiograph suggested that each canal was independent



**Figure 1:** Preoperative radiograph showing periapical radiolucent areas.



**Figure 2:** Radiograph for determination of working length shows only 1 canal in each tooth.



**Figure 3:** A second radiograph for determination of working length shows 6 distinct and independent canals.



**Figure 4:** Postoperative radiograph of the left central incisor and the left lateral incisor.



**Figure 5:** Postoperative radiograph of the right central incisor.

throughout its course and possessed its own apex opening (Fig. 3).

At the third visit, all of the canals were filled with a thermoplasticized gutta-percha. Temporary zinc oxyde-eugenol (ZOE) restorations were placed, and permanent restorations were advised. Postoperative radiographs showed 2 separate canals and 2 separate foramina in each tooth (Figs. 4 and 5).

1 patient. The morphological pattern of separate apical terminations for 2 canals in multiple mandibular incisors, as manifested in this case, is very rare. ♦



*Dr. Hwang is a graduate student in the department of conservative dentistry at Wonkwang University, College of Dentistry, Iksan, Cheonbuk, South Korea.*



*Dr. Min is instructor in the department of conservative dentistry at Wonkwang University, College of Dentistry, Iksan, Cheonbuk, South Korea.*

*Correspondence to: Dr. Kyung san Min, Wonkwang University, College of Dentistry, 344-2 Sinyong, Iksan, Choebuk, South Korea. 570-749. E-mail : mksdd@wonkwang.ac.kr.*

*The authors have no declared financial interests in any company manufacturing the types of products mentioned in this article.*

## Discussion

The main objective of root canal treatment is thorough mechanical and chemical cleansing of the entire pulp space and complete obturation with an inert filling material.<sup>7</sup> According to the endodontic literature, mandibular incisors with 2 canals are not unusual.<sup>1-4</sup> In most cases in which 2 canals are present, however, they merge into a single canal short of the apex. The occurrence of 2 separate foramina in mandibular incisors with 2 canals has been described in only a limited number of reports.<sup>8</sup> Careful interpretation of the radiographic features is essential to ensure that additional root canals are not overlooked. This may necessitate imaging the tooth from a variety of angles so that the root canals may be distinguished in the resulting films. In addition, if an obvious canal ends abruptly, the clinician should be suspicious that there are in fact 2 canals.<sup>9</sup> Nevertheless, manual exploration of the root canal system with an endodontic file or explorer is a reliable way to identify the exact configuration of the root canal, especially the number of foramina. This report has described the root canal treatment of 3 mandibular incisors, each with 2 separate canals having their own foramina, in

## References

1. Vertucci FJ. Root canal anatomy of the mandibular anterior teeth. *J Am Dent Assoc* 1974; 89(2):369-71.
2. Bellizzi R, Hartwell G. Clinical investigation of in vivo endodontically treated mandibular anterior teeth *J Endod* 1983; 9(6):246-8.
3. Rankine-Willson RW, Henry P. The bifurcated root canals in lower anterior teeth. *J Am Dent Assoc* 1965; 70:1162-5.
4. Benjamin KA, Dowson J. Incidence of two root canals in human mandibular incisor teeth. *Oral Surg Oral Med Oral Pathol* 1974; 38(1):122-6.
5. Miyashita M, Kasahara E, Yasuda E, Yamamoto A, Sekizawa T. Root canal system of the mandibular incisor. *J Endod* 1997; 23(8):479-84.
6. Mauger MJ, Schindler WG, Walker WA 3rd. An evaluation of canal morphology at different levels of root resection in mandibular incisors. *J Endod* 1998; 24(9):607-9.
7. Christie WH, Peikoff MD, Acheson DW. Endodontic treatment of two maxillary lateral incisors with anomalous root formation. *J Endod* 1981; 7(11):528-34.
8. Funato A, Funato H, Matsumoto K. Mandibular central incisor with two root canals. *Endod Dent Traumatol* 1998; 14(6):285-6.
9. Cohen S, Burns RC. Pathways of the pulp. 8th ed. St. Louis: Mosby Year Book Inc.; 2002. p. 202.