

# Treatment of Self-Inflicted Oral Trauma in a Comatose Patient: A Case Report

**Taira Kobayashi, DDS, PhD; Henry Ghanem, BDS, LDS, MS;**  
**Koji Umezawa, DDS, PhD; Junichi Mega, DDS, PhD;**  
**Misao Kawara, DDS, PhD; Jocelyne S. Feine, DDS, HDR**

## Contact Author

**Dr. Ghanem**  
E-mail:  
henry.ghanem@mcgill.ca



## ABSTRACT

Trauma to oral soft tissues in comatose patients may be more widespread than reported, as no extensive study of this problem has been conducted. Various appliances for the prevention of self-inflicted injuries to oral tissues particularly in children and the physically and mentally challenged have been documented, but there is little information on their use in adult comatose patients. Because comatose patients lack cerebral control of the masticatory cycle, they can easily injure themselves. Although it is not uncommon for patients with a decreased level of consciousness and in need of intensive care to be restrained to prevent injury due to involuntary movement of the limbs, head and neck restraint is often difficult and may be dangerous to the patient. This case report presents a simple solution to the problem of self-inflicted trauma to oral tissues.

**MeSH Key Words:** coma; lip/injuries; mouth protectors; self mutilation/prevention & control

© J Can Dent Assoc 2005; 71(9):661–4  
This article has been peer reviewed.

A team effort is needed in the management of the comatose patient with self-inflicted oral trauma. On occasion, the private dental practitioner may be asked to provide clinical assistance to the team. Prevalence of trauma to oral tissues in the comatose patient is not well documented, but when it occurs it is quite destructive and distressful for both patient and family members. Most reported cases have been in the pediatric or special care journals,<sup>1–22</sup> with few reported for the adult decerebrate patient.

The cause of self-injurious behaviour and clinical symptoms varies and the literature is full of examples, especially in children with cerebral palsy,<sup>1,16,18</sup> severe neuro-disability,<sup>23</sup> Tourette's syndrome,<sup>3,19</sup> Lesch-Nyhan syndrome,<sup>4,5,9</sup> Chiari Type II malformations,<sup>7</sup> congenital insensitivity to pain,<sup>6,8,12</sup> various psychiatric disorders and encephalitis in infancy.<sup>11</sup>

Lack of control over the masticatory cycle in the comatose adult patient may sometimes result in neuropathologic chewing,<sup>21,24</sup> which

in turn may be the result of severe brain damage due to a closed head injury, hypoxia and septic shock.<sup>13</sup> Management of these cases will vary according to the individual's medical history, frequency and severity of the injury and whether treatment is for the long or short term.

Numerous appliances and techniques have been advocated: occlusal bite planes, ratchet mouth props, padded tongue blades and intermaxillary fixation, and mandibular-cast silver caps with acrylic bite-blocks.<sup>13</sup> There are no standards, but it is universally accepted that a removable device is desirable for long-term use in comatose patients. Difficulties that may be encountered with respect to the design of an appliance include lack of cooperation from the patient, inability to gain access to the oral cavity for a proper examination and to take impressions, and the education of caregivers and family members in maintenance of the appliance and proper oral hygiene procedures for the patient.

An appropriate appliance must be simple to make and be well retained and easily serviced. It must also satisfy the following criteria<sup>25</sup>:


- deflect the tissues most likely to be damaged by involuntary movements of the mandible from the occlusal table
- permit a full range of mandibular movement
- allow for daily oral care
- withstand breakage and displacement forces over an indefinite period
- allow healing of traumatized tissues
- be easily fabricated and placed without risk to the patient.

The following case report describes the use of a simple device in an adult comatose patient. The material used is easily layered or repaired using heat alone without the need for organic solvents.

### Case Report

The dental service of the Nihon University School of Dentistry at Matsudo was asked to examine and recommend treatment for a 56-year-old male patient who was intubated in hospital after suffering a subarachnoid hemorrhage caused by the rupture of a cerebral aneurysm. He had sustained self-induced lacerations to his lower lip. The patient underwent a ventriculoperitoneal shunt insertion and nutrition was provided by means of a nasogastric tube following tracheotomy. According to the Japan coma scale, his level of consciousness was III-200 (Box 1). At this level of consciousness, the patient could involuntarily move his arms and legs and frown on painful stimulus, but could not respond to requests, for example, to open his mouth. His medical condition was stable. Hospital staff attempted to control his involuntary masticatory movements by inserting gauze and tongue spatulas between his teeth but this failed (Fig. 1). The patient had been examined previously by another dentist, who had suggested extraction of all lower mandibular incisors to prevent injury. This recommendation is a possible solution, but only as a last resort when all other options have failed.

A thorough examination of the patient was difficult due to the restricted opening of his jaws, but he appeared to have a significant overjet and overbite, suggesting a Class II, Division 1 occlusion. In addition, there was a severe self-inflicted traumatic lesion on his lower lip mainly on the left side (Fig. 2) and a transient ruminatory chewing cycle was observed. This suggested that the lower lip was trapped between the maxillary and mandibular



**Figure 1:** Patient with gauze wrapped around spatula between teeth.



**Figure 2:** Laceration on lower lip.



**Figure 3:** Polyolefin mouthguard.



**Figure 4:** Mouthguard with handle.

### Box 1 Japan coma scale; 3-3-9 system

#### 1. Wakes even without stimulus

- 1 Almost lucid, but consciousness is not as clear as it ought to be
- 2 Disoriented
- 3 Unable to recall name or birthday

#### 2. Wakes with stimulus, but falls asleep when stimulus ceases

- 10 Opens eyes readily when called
- 20 Opens eyes in response to loud calling or shaking
- 30 Opens eyes narrowly with repeated calling or painful stimulus

#### 3. Does not wake even with stimulus

- 100 Performs motions such as brushing off painful stimulus
- 200 Moves arms and legs or frowns with painful stimulus
- 300 No response to painful stimulus

anterior teeth and was acting as a bolus, thus initiating a pathologic chewing cycle. The ulcer created by this action was sizeable; the tissue was edematous and bleeding. All maxillary and mandibular teeth were present except for teeth 36 and 46, which had been replaced with two 3-unit fixed partial dentures. Observed laceration of the lower lip was due to adduction of the lip as a result of sucking and bruxism.



**Figure 5:** Mouthguard in patient's mouth.



**Figure 6:** Initial resolution of wound 3 days after placement of mouthguard.



**Figure 7:** Two weeks after insertion, fibrotic scar is visible.



**Figure 8:** Complete resolution of wound.

Because of the difficulty encountered in trying to open the jaw due to trismus, we decided to make a simple, easily serviceable appliance that would satisfy the criteria outlined by Hanson.<sup>25</sup> With great difficulty, a gag was used to open the mouth and the opening was secured on the right side with a medium-sized McKesson rubber bite-block (McKesson Mouth Props, Hu-Friedy, Chicago, Ill.).

A unilateral impression was made using a "rim lock" stock tray and irreversible hydrocolloid impression material. A conventional mouthguard was fabricated using a polyolefin sheet (MG21, Molten Medical Inc., Tokyo, Japan) with adequate coverage of teeth 33–47 to ensure good retention (Fig. 3), but not overly extended to avoid impingement on soft tissue and muscle attachment. A long flexible plastic handle made of the same material was attached to the anterior portion of the mouthguard (Fig. 4); this could be secured to the patient's hospital gown with a safety pin to prevent accidental loss or possible inhalation of the appliance should it become dislodged. The labial portion of the appliance was slightly thickened (8 mm) by layering to prevent the lips from becoming trapped between the anterior teeth (Fig. 5). The nursing staff were asked to keep the appliance in the mouth at all times except when it was removed for daily oral hygiene procedures.

## Results

Within 3 to 4 days after insertion of the appliance, the nursing staff noted that serious clenching of the jaw had ceased and that it was possible to manipulate the jaw with a tapping motion. Sucking and adduction of the lips had ceased and initial resolution of the wound was observed (Fig. 6). By 14 days after insertion, healing of the wound was noted, with the appearance of a fibrotic scar (Fig. 7). At 4 weeks postinsertion, the lesion had completely resolved.

Although the appliance was removed after healing (Fig. 8), the family asked that it be replaced as a preventive measure in the long term, because they noted that periodically the patient adducted the lips between the anterior teeth. It was decided to comply with the wishes of the family with the proviso that oral hygiene and maintenance would be carried out; a full coverage mandibular mouthguard would be constructed at a later date to avoid any occlusal problems that might arise from wearing a unilateral mouthguard for an extended period; and the patient would be seen by a dentist every 3 months.

## Discussion

Trauma to oral soft tissue in adult comatose patients may be more widespread than documented and, although no extensive study has been conducted on the subject, many authors have suggested the fabrication of appliances to prevent self-inflicted trauma, particularly in children.

Mastication and masticatory movements have been described as the result of a complex set of actions that are both voluntary and automatic; these movements involve the coordination of the muscles of mastication operating in a learned pattern to prepare and break down a bolus of food for swallowing. Learning and control of these movements is under the influence of the central cortex, reticular formation and the extra-pyramidal system.<sup>21</sup> One characteristic of lack of cerebral control is lack of coordination of the masticatory movement, which in turn may lead to self-inflicted trauma to oral tissues. Bruxism seems to appear at different levels of unconsciousness and disappears only after a significant improvement in the level of consciousness.<sup>22</sup> Myostatic masticatory reflex may be initiated in a comatose patient if the lips or tongue are trapped between the teeth, mimicking the placement of a bolus of food on the occlusal surfaces of the teeth.<sup>26</sup>

Management of these patients should be immediate, using an easily fabricated removable appliance in the short term, and a long-term solution should be sought as level of consciousness improves. Ngan and Nelson<sup>21</sup> state that the design of any appliance depends on the prognosis of the patient, neurologic status and the severity of chewing or bruxing. The practice has been to construct either a full-coverage mouthguard or one covering the anterior sextant using traditional materials; however, these materials lack proper adaptation to the dentition, hence are not adequately retained. Dislodgement of such an appliance may cause even more serious complications if it is inhaled. In this case, a simple cross-arch appliance made of polyolefin was introduced. This mouthguard material is odorless, has good visco-elastic properties and high impact resistance and is easy to laminate and repair by layering using heat rather than adhesives. Because the material is also lightweight and highly resistant to tearing, we were able to fabricate a long handle that could be secured to the patient's garment making it easy for caregivers to remove and replace the device during oral hygiene procedures. The cross-arch design, which covered two-thirds of the mandibular occlusal surface, was successful in resolving the problem in the short term. In the long term, a full arch stent would be more appropriate to prevent an imbalance to the occlusion and inadvertent overeruption of teeth not covered by a partial appliance. ♦

## THE AUTHORS



*Dr. Kobayashi is a visiting professor in the faculty of dentistry at McGill University, Montreal, Quebec.*



*Dr. Ghanem is a doctoral candidate in the faculty of dentistry at McGill University, Montreal, Quebec.*



*Dr. Umezawa is an assistant professor in the department of dentistry for the disabled, Nihon University School of Dentistry at Matsudo, Japan.*



*Dr. Mega is a professor in the department of dentistry for the disabled, Nihon University School of Dentistry at Matsudo, Japan.*



*Dr. Kawara is a professor in the department of comprehensive clinical dentistry, Nihon University School of Dentistry at Matsudo, Japan.*



*Dr. Feine is a professor in the faculty of dentistry at McGill University, Montreal, Quebec.*

*Correspondence to:* Dr. Henry Ghanem, Faculty of Dentistry, McGill University, 3640 University St., Montreal, QC H3A 2B2. E-mail: henry.ghanem@mcgill.ca.

*The authors have no declared financial interests in any company manufacturing the types of products mentioned in this article.*

## References

1. Yasui EM, Kimura RK, Kawamura A, Akiyama S, Morisaki I. A modified oral screen appliance to prevent self-inflicted oral trauma in an infant with cerebral palsy: a case report. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2004; 97(4):471-5.
2. Silva DR, da Fonseca MA. Self-injurious behavior as a challenge for the dental practice: a case report. *Pediatr Dent* 2003; 25(1):62-6.
3. Simoyama T, Horie N, Kato T, Nasu D, Kaneko T. Tourette's syndrome with rapid deterioration by self-mutilation of the upper lip. *J Clin Pediatr Dent* 2003; 27(2):177-85.
4. Fardi K, Topouzelia N, Kotsanos N. Lesch-Nyham syndrome: a preventive approach to self-mutilation. *Int J Paediatr Dent* 2003; 13(1):51-6.
5. Cusumano FJ, Penna KJ, Panossian G. Prevention of self-mutilation in patients with Lesch-Nyhan syndrome: review of literature. *ASDC J Dent Child* 2001; 68(3):175-8.
6. Erdem TL, Ozcan I, Ilguy D, Sirin S. Hereditary sensory and autonomic neuropathy: review and a case report with dental implications. *J Oral Rehabil* 2000; 27(2):180-3.
7. Nurko C, Errington BD, Ben Taylor W, Henry R. Lip biting in a patient with Chiari type II malformation: case report. *Pediatr Dent* 1999; 21(3):209-12.
8. Littlewood SJ, Mitchell L. The dental problem and management of a patient suffering from congenital insensitivity to pain. *Int J Paediatr Dent* 1998; 8(1):47-50.
9. Rashid N, Yusuf H. Oral self-mutilation by a 17-month-old child with Lesch-Nyhan syndrome. *Int J Paediatr Dent* 1997; 7(2):115-7.
10. Saemundsson SR, Roberts MW. Oral self-injurious behavior in the developmentally disabled: review and a case. *ASDC J Dent Child* 1997; 64(3):205-9, 228.
11. Cehreli ZC, Olmez S. The use of a special mouthguard in the management of oral injury self-inflicted by a 4-year-old child. *Int J Paediatr Dent* 1996; 6(4):277-81.
12. Rasmussen P. The congenital insensitivity-to-pain syndrome (analgesia congenital): report of a case. *Int J Paediatr Dent* 1996; 6(2):117-22.
13. Hallett KB. Neuropathological chewing: a dental management protocol and treatment appliances for pediatric patients. *Spec Care Dent* 1994; 14(2):61-4.
14. Sheller B. Self-inflicted oral trauma: report of case. *Spec Care Dent* 1992; 12(1):28-9.
15. Finger ST, Duperon DF. The management of self-inflicted oral trauma secondary to encephalitis: a clinical report. *ASDC J Dent Child* 1991; 58(1):60-3.
16. Sonnenberg EM. Treatment of self-induced trauma in a patient with cerebral palsy. *Spec Care Dent* 1990; 10(3):89-90.
17. Croglio DP, Thines TJ, Fleischer MS, Anders PL. Self-inflicted oral trauma. *Spec Care Dent* 1990; 10(2):58-61.
18. Rover BC, Morgano SM. Prevention of self-inflicted trauma: dental intervention to prevent chronic lip chewing by a patient with a diagnosis of progressive bulbar palsy. *Spec Care Dent* 1998; 8(1):37-9.
19. Lowe O. Tourette's syndrome: management of oral complications. *ASDC J Dent Child* 1986; 53(6):456-60.
20. Peters TE, Blair AE, Freeman RG. Prevention of self-inflicted trauma in comatose patients. *Oral Surg Oral Med Oral Pathol* 1984; 57(4):367-70.
21. Ngan PW, Nelson LP. Neuropathologic chewing in comatose children. *Pediatr Dent* 1985; 7(4):302-6.
22. Pratap-Chand R, Gourie-Devi M. Bruxism: its significance in coma. *Clin Neurol Neurosurg* 1985; 87(2):113-7.
23. Millwood J, Fiske J. Lip-biting in patients with profound neuro-disability. *Dent Update* 2001; 28(2):105-8.
24. Guyton AC. Textbook of medical physiology. 3rd ed. Philadelphia: WB Saunders Co; 1966. p. 879.
25. Hanson GE, Ogle RG, Giron L. A tongue stent for prevention of oral trauma in the comatose patient. *Critical Care Med* 1975; 3(5):200-3.
26. Jackson MJ. The use of tongue-depressing stents for neuropathologic chewing. *J Prosthet Dent* 1978; 40(3):309-11.